



## **Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection**

Downloaded from <https://aidsinfo.nih.gov/guidelines> on 8/31/2020

Visit the AIDSinfo website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <https://aidsinfo.nih.gov/e-news>.

**Table 15a. Antiretroviral Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System Toxicity** (Last updated April 14, 2020; last reviewed April 14, 2020) (page 1 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
<b>Global CNS Depression</b>	LPV/r oral solution (contains both ethanol and propylene glycol as excipients)	<b>Onset:</b> <ul style="list-style-type: none"> <li>1–6 days after starting LPV/r</li> </ul> <b>Presentation</b> <i>Neonates/Premature Infants:</i> <ul style="list-style-type: none"> <li>Global CNS depression (e.g., abnormal EEG, altered state of consciousness, somnolence)</li> </ul>	Unknown; rare case reports have been published	Prematurity  Low birth weight  Aged <14 days (whether birth was premature or term)	Avoid use of LPV/r until a postmenstrual age of 42 weeks and a postnatal age of ≥14 days unless no other alternatives are available, <a href="#">see Lopinavir/Ritonavir</a> .	Discontinue LPV/r; symptoms should resolve in 1–5 days.  If needed, reintroduction of LPV/r can be considered once the patient is outside the vulnerable period (i.e., postmenstrual age of 42 weeks and a postnatal age ≥14 days).
<b>Neuropsychiatric Symptoms and Other CNS Manifestations</b>	EFV	<b>Onset:</b> <ul style="list-style-type: none"> <li>For many symptoms, onset is 1–2 days after starting EFV.</li> <li>Many symptoms subside or diminish by 2–4 weeks, but symptoms may persist in a significant proportion of patients.</li> </ul> <b>Presentation (May Include One or More of the Following)</b> <i>Neuropsychiatric Symptoms:</i> <ul style="list-style-type: none"> <li>Abnormal dreams</li> <li>Psychosis</li> <li>Suicidal ideation or attempted/completed suicide</li> </ul> <i>Other CNS Manifestations:</i> <ul style="list-style-type: none"> <li>Dizziness</li> <li>Somnolence</li> <li>Insomnia or poor sleep quality</li> <li>Impaired concentration</li> <li>Seizures (including absence seizures)</li> <li>Cerebellar dysfunction (e.g., tremor, dysmetria, ataxia)</li> </ul> <b>Note:</b> CNS side effects (e.g., impaired concentration, abnormal dreams, sleep disturbances) may be more difficult to assess in children.	Variable, depending on age, symptoms, and assessment method  <b>Children:</b> <ul style="list-style-type: none"> <li>24% of patients experienced any EFV-related CNS manifestation in one case series, with 18% of participants requiring drug discontinuation.</li> <li>Five of 45 participants (11%) experienced new-onset seizures in one study of children aged &lt;36 months; two of these participants had alternative causes for seizures.</li> <li>Cases of cerebellar dysfunction have been reported in children with very high EFV plasma levels.</li> </ul> <b>Adults:</b> <ul style="list-style-type: none"> <li>30% incidence for any CNS manifestations of any severity.</li> <li>6% incidence for EFV-related, severe CNS manifestations, including suicidality. However, evidence is conflicting about whether EFV use increases the incidence of suicidality.</li> <li>One case series reported 20 women with ataxia that resolved upon EFV discontinuation, but frequency was not reported.</li> </ul>	Insomnia is associated with elevated EFV trough concentration (≥4 mcg/mL)  CYP2B6 polymorphisms that decrease EFV metabolism and cause increased EFV serum concentrations (CYP2B6 516 T/T genotype or co-carriage of CYP2B6 516 G/T and 983 T/C variants)  History of psychiatric illness or use of psychoactive drugs	Administer EFV on an empty stomach, preferably at bedtime.  Prescreen for psychiatric illness; avoid use in the presence of psychiatric illness, including depression or suicidal thoughts. Avoid concomitant use of psychoactive drugs.  Consider using TDM in children with mild or moderate EFV-associated toxicities.	If symptoms are excessive or persistent, obtain EFV trough concentration. If EFV trough concentration is >4 mcg/mL and/or symptoms are severe, strongly consider drug substitution if a suitable alternative exists.  Alternatively, consider dose reduction with repeat TDM and dose adjustment (with input from an expert pharmacologist).

**Table 15a. Antiretroviral Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System Toxicity** (Last updated April 14, 2020; last reviewed April 14, 2020) (page 2 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
<b>Neuropsychiatric Symptoms and Other CNS Manifestations,</b> continued	RPV	<b>Onset:</b> <ul style="list-style-type: none"> <li>Most symptoms occur in the first 4–8 weeks of treatment.</li> </ul> <b>Presentation</b> <i>Neuropsychiatric Symptoms:</i> <ul style="list-style-type: none"> <li>Depressive disorders</li> <li>Suicidal ideation</li> <li>Abnormal dreams/nightmares</li> </ul> <i>Other CNS Manifestations:</i> <ul style="list-style-type: none"> <li>Headache</li> <li>Dizziness</li> <li>Insomnia</li> <li>Somnolence</li> </ul>	<b>Adults:</b> <ul style="list-style-type: none"> <li>CNS/neuro-psychiatric adverse events of all severity grades were reported in 43% of patients at 96 weeks (most were Grade 1). Depressive disorders of all severity grades were reported in 9% of patients; 1% of patients discontinued RPV due to severe depressive disorders.</li> </ul> <b>Children:</b> <ul style="list-style-type: none"> <li>Depressive disorders of all severity grades were reported in 19.4% of pediatric patients aged 12–17 years. Severe depressive disorders were reported in 5.6% of patients, including one suicide attempt.</li> <li>Somnolence was reported in five of 36 children (14%).</li> </ul>	History of neuropsychiatric illness	Monitor carefully for depressive disorders and other CNS symptoms.	Consider drug substitution in cases of severe symptoms.
	RAL	<b>Onset:</b> <ul style="list-style-type: none"> <li>As early as 3–4 days after starting RAL</li> </ul> <b>Presentation:</b> <ul style="list-style-type: none"> <li>Increased psychomotor activity</li> <li>Headaches</li> <li>Insomnia</li> <li>Depression</li> <li>Cerebellar dysfunction (e.g., tremor, dysarthria, ataxia)</li> </ul>	<b>Children:</b> <ul style="list-style-type: none"> <li>Increased psychomotor activity was reported in one child.</li> </ul> <b>Adults:</b> <ul style="list-style-type: none"> <li>Headache</li> <li>Insomnia (&lt;5% in adult trials)</li> <li>Rare case reports of cerebellar dysfunction in adults</li> </ul>	Elevated RAL concentrations  Co-treatment with TDF, a PPI, or inhibitors of UGT1A1  Prior history of insomnia or depression	Prescreen for psychiatric symptoms.  Monitor carefully for CNS symptoms.  Use with caution in the presence of drugs that increase RAL concentration.	Consider drug substitution (RAL or coadministered drug) in cases of severe insomnia or other neuropsychiatric symptoms.

**Table 15a. Antiretroviral Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System Toxicity** (Last updated April 14, 2020; last reviewed April 14, 2020) (page 3 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
<b>Neuropsychiatric Symptoms and Other CNS Manifestations,</b> continued	DTG	<p><b>Onset:</b></p> <ul style="list-style-type: none"> <li>• 7–30 days after starting DTG</li> </ul> <p><b>Presentation</b></p> <p><i>Neuropsychiatric Symptoms:</i></p> <ul style="list-style-type: none"> <li>• Depression or exacerbation of preexisting depression</li> <li>• Anxiety</li> <li>• Suicidal ideation or attempted/completed suicide</li> <li>• <b>Drowsiness</b></li> <li>• <b>Neurocognitive deficits (lower total competence and school performance)</b></li> </ul> <p><i>Other CNS Manifestations (Generally Mild):</i></p> <ul style="list-style-type: none"> <li>• Sleep disturbances</li> <li>• Dizziness</li> <li>• Headache</li> </ul>	<p><b>Children:</b></p> <ul style="list-style-type: none"> <li>• <b>In a retrospective cohort analysis, neuropsychiatric events that resulted in discontinuation occurred in two of 29 (6.8%) children who initiated DTG.</b></li> </ul> <p><b>Adults:</b></p> <ul style="list-style-type: none"> <li>• <b>2.7% of the neuropsychiatric AEs reported in a large prospective cohort resulted in treatment discontinuation.</b></li> <li>• Higher frequency of neuropsychiatric symptoms reported with DTG than with other INSTIs. A class effect has been suggested.</li> </ul>	<p>Pre-existing depression or other psychiatric illness</p> <p><b>History of ARV-related neuropsychiatric symptoms</b></p> <p>Higher frequency of neuropsychiatric symptoms reported when DTG is coadministered with ABC; however, evidence is conflicting.</p> <p>UGT1A1*6 and/or *28 polymorphism (reported in patients of Asian descent)</p>	<p>Use with caution in the presence of psychiatric illness, especially in patients with depression <b>or a history of ARV-related neuropsychiatric symptoms.</b></p> <p>Consider morning dosing of DTG.</p>	<p>For persistent or severe neuropsychiatric symptoms, consider discontinuing DTG if a suitable alternative exists.</p> <p>For mild symptoms, continue DTG and counsel patient that symptoms will likely resolve with time.</p>

**Table 15a. Antiretroviral Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System Toxicity** (Last updated April 14, 2020; last reviewed April 14, 2020) (page 4 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Neuropsychiatric Symptoms and Other CNS Manifestations, continued	BIC	<p><b>Onset:</b></p> <ul style="list-style-type: none"> <li>1–63 days after starting BIC (as late as 233 days for schizoaffective disorders)</li> </ul> <p><b>Presentation</b></p> <p><i>Neuropsychiatric Symptoms:</i></p> <ul style="list-style-type: none"> <li>Depression or exacerbation of pre-existing depression</li> <li>Suicidal ideation or attempted suicide</li> <li>Schizoaffective disorders</li> <li>Anxiety</li> </ul> <p><i>Other CNS Manifestations (Generally Mild):</i></p> <ul style="list-style-type: none"> <li>Abnormal dreams</li> <li>Dizziness</li> <li>Insomnia</li> </ul>	<p>Data in children and adults come mostly from clinical trials. Overall, the frequency of neuropsychiatric events in BIC and DTG comparator arms appeared similar in adult clinical trials.</p> <p><b>Children:</b></p> <ul style="list-style-type: none"> <li>1 child (1%) had Grade 2 insomnia and anxiety that led to drug discontinuation.</li> </ul> <p><b>Adults:</b></p> <ul style="list-style-type: none"> <li>Abnormal dreams, dizziness, and insomnia occurred in 1% to 5% of adults.</li> <li>Suicidal ideation, suicide attempts, schizoaffective disorders, and depression occurred in &lt;1% of adults.</li> </ul>	<p>Pre-existing depression or other psychiatric conditions</p> <p>History of ARV-related neuropsychiatric symptoms</p>	<p>Use with caution in the presence of psychiatric conditions, or in patients with a history of ARV-related neuropsychiatric symptoms.</p>	<p>For persistent or severe neuropsychiatric symptoms, consider discontinuing BIC if a suitable alternative exists.</p> <p>For mild symptoms, continue BIC and counsel patient that symptoms will likely resolve with time.</p>

**Key:** ABC = abacavir; ARV = antiretroviral; BIC = bictegravir; CNS = central nervous system; CYP = cytochrome P; DTG = dolutegravir; EEG = electroencephalogram; EFV = efavirenz; INSTI = integrase strand transfer inhibitor; LPV/r = lopinavir/ritonavir; PPI = proton pump inhibitor; RAL = raltegravir; RPV = rilpivirine; TDF = tenofovir disoproxil fumarate; TDM = therapeutic drug monitoring; UGT = uridine diphosphate-glucuronosyltransferase

## References

1. Arenas-Pinto A, Grund B, Sharma S, et al. Increased risk of suicidal behaviour with use of efavirenz: results from the START trial. Presented at: 21st International AIDS Conference. 2016. Durban, South Africa.
2. Bracchi M, Pagani N, et al. Multicentre open-label pilot study of switching from efavirenz to dolutegravir for central nervous system (CNS) toxicity. Abstract # P209. Presented at: HIV Drug Therapy. 2016. Glasgow, Scotland. Available at: <http://hivglasgow.org/2016-abstracts>.
3. Briand C, Dollfus C, Faye A, et al. Efficacy and tolerance of dolutegravir-based combined ART in perinatally HIV-1-infected adolescents: a French multicentre retrospective study. *J Antimicrob Chemother*. 2017;72(3):837-843. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27999017>.
4. Cabrera Figueroa S, Fernandez de Gatta M, Hernandez Garcia L, et al. The convergence of therapeutic drug monitoring and pharmacogenetic testing to optimize efavirenz therapy. *Ther Drug Monit*. 2010;32(5):579-585. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20720517>.
5. Cattaneo D, Ripamonti D, Baldelli S, Cozzi V, Conti F, Clementi E. Exposure-related effects of atazanavir on the pharmacokinetics of raltegravir in HIV-1-infected

*Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection*

K-9

- patients. *Ther Drug Monit*. 2010;32(6):782-786. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20926993>.
6. Cohen CJ, Molina JM, Cassetti I, et al. Week 96 efficacy and safety of rilpivirine in treatment-naïve, HIV-1 patients in two Phase III randomized trials. *AIDS*. 2013;27(6):939-950. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23211772>.
  7. de Boer MG, van den Berk GE, van Holten N, et al. Intolerance of dolutegravir-containing combination antiretroviral therapy regimens in real-life clinical practice. *AIDS*. 2016;30(18):2831-2834. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27824625>.
  8. Fernandez C, Michie K, et al. Adverse events and discontinuation of dolutegravir-based therapy in naïve and experienced HIV patients: tertiary HIV centre experience. Abstract # P212. Presented at: HIV Drug Therapy. 2016. Glasgow, Scotland. Available at: <http://hivglasgow.org/2016-abstracts>.
  9. Fettiplace A, Stainsby C, Winston A, et al. Psychiatric symptoms in patients receiving dolutegravir. *J Acquir Immune Defic Syndr*. 2017;74(4):423-431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27984559>.
  10. Rilpivirine [package insert]. Food and Drug Administration. 2018. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/202022s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/202022s011lbl.pdf).
  11. Food and Drug Administration. FDA drug safety communication: serious health problems seen in premature babies given kaletra (lopinavir/ritonavir) oral solution. 2011. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm246002.htm>.
  12. Ford N, Shubber Z, Pozniak A, et al. Comparative safety and neuropsychiatric adverse events associated with efavirenz use in first-line antiretroviral therapy: A systematic review and meta-analysis of randomized trials. *J Acquir Immune Defic Syndr*. 2015;69(4):422-429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25850607>.
  13. García-Navarro C, Jiménez de Ory S, Navarro Gómez ML, et al. Sleep disturbances in a cohort of HIV-infected children and adolescents on antiretroviral treatment. Presented at: 22nd Conference on Retroviruses and Opportunistic Infections. 2015. Seattle, WA.
  14. Group ES, Carey D, Puls R, et al. Efficacy and safety of efavirenz 400 mg daily versus 600 mg daily: 96-week data from the randomised, double-blind, placebo-controlled, non-inferiority ENCORE1 study. *Lancet Infect Dis*. 2015;15(7):793-802. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25877963>.
  15. Haas DW, Ribaud HJ, Kim RB, et al. Pharmacogenetics of efavirenz and central nervous system side effects: an adult AIDS clinical trials group study. *AIDS*. 2004;18(18):2391-2400. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15622315>.
  16. Hauptfleisch MP, Moore DP, Rodda JL. Efavirenz as a cause of ataxia in children. *S Afr Med J*. 2015;105(11):897-898. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26632310>.
  17. Kheloufi F, Allemann J, Mokhtari S, Default A. Psychiatric disorders after starting dolutegravir: report of four cases. *AIDS*. 2015;29(13):1723-1725. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26372287>.
  18. Leutscher PD, Stecher C, Storgaard M, Larsen CS. Discontinuation of efavirenz therapy in HIV patients due to neuropsychiatric adverse effects. *Scand J Infect Dis*. 2013;45(8):645-651. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23427878>.
  19. Lombaard J, Bunupuradah T. Week 48 safety and efficacy of a rilpivirine (TMC278)-based regimen in HIV-infected treatment-naïve adolescents: PAINT Phase II trial. Presented at: 7th International Workshop on HIV Pediatrics. 2015. Vancouver, Canada.
  20. Madeddu G, Menzaghi B, Ricci E, et al. Raltegravir central nervous system tolerability in clinical practice: results from a multicenter observational study. *AIDS*. 2012;26(18):2412-2415. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23032413>.



21. Mills A, Garner W, Pozniak A, et al. Patient-reported symptoms over 48 weeks in a randomized, open-label, Phase IIIb non-inferiority trial of adults with HIV switching to co-formulated elvitegravir, cobicistat, emtricitabine, and tenofovir DF versus continuation of non-nucleoside reverse transcriptase inhibitor with emtricitabine and tenofovir DF. *Patient*. 2015;8(4):359-371. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26045359>.
22. Mollan KR, Smurzynski M, Eron JJ, et al. Association between efavirenz as initial therapy for HIV-1 infection and increased risk for suicidal ideation or attempted or completed suicide: an analysis of trial data. *Ann Intern Med*. 2014;161(1):1-10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24979445>.
23. Mollan KR, Tierney C, Eron JJ, et al. Composite CYP2B6/CYP2A6 genotype and risk for suicidality among HIV-infected individuals randomly assigned to initiate efavirenz-containing regimens in AIDS Clinical Trials Group studies. Presented at: 21st International AIDS Conference. 2016. Durban, South Africa.
24. Nachman S, et al. IMPAACT P1066: raltegravir (RAL) safety and efficacy in HIV infected (+) youth two to 18 years of age through week 48. Abstract no. TUAB0205. Presented at: 19th International AIDS Conference. 2012. Washington, DC.
25. Napoli AA, Wood JJ, Coumbis JJ, Soitkar AM, Seekins DW, Tilson HH. No evident association between efavirenz use and suicidality was identified from a disproportionality analysis using the FAERS database. *J Int AIDS Soc*. 2014;17:19214. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25192857>.
26. Nkhoma ET, Coumbis J, Farr AM, et al. No evidence of an association between efavirenz exposure and suicidality among HIV patients initiating antiretroviral therapy in a retrospective cohort study of real world data. *Medicine*. 2016;95(3):e2480. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26817882>.
27. Pinillos F, Dandara C, Swart M, et al. Case report: severe central nervous system manifestations associated with aberrant efavirenz metabolism in children: the role of CYP2B6 genetic variation. *BMC Infect Dis*. 2016;16:56. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26831894>.
28. Puthanakit T, Tanpaiboon P, Aupibul L, Cressey TR, Sirisanthana V. Plasma efavirenz concentrations and the association with CYP2B6-516G >T polymorphism in HIV-infected Thai children. *Antivir Ther*. 2009;14(3):315-320. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19474465>.
29. Quercia R, Roberts J. Psychiatric adverse events from the DTG ART-naïve Phase 3 clinical trials. Abstract # P210. Presented at: HIV Drug Therapy. 2016. Glasgow, Scotland. Available at: <http://hivglasgow.org/2016-abstracts>.
30. Rakhmanina NY, van den Anker JN, Soldin SJ, van Schaik RH, Mordwinkin N, Neely MN. Can therapeutic drug monitoring improve pharmacotherapy of HIV infection in adolescents? *Ther Drug Monit*. 2010;32(3):273-281. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20445485>.
31. Reiss KA, Bailey JR, Pham PA, Gallant JE. Raltegravir-induced cerebellar ataxia. *AIDS*. 2010;24(17):2757. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20980871>.
32. Shubber Z, Calmy A, Andrieux-Meyer I, et al. Adverse events associated with nevirapine and efavirenz-based first-line antiretroviral therapy: a systematic review and meta-analysis. *AIDS*. 2013;27(9):1403-1412. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23343913>.
33. Smith C, Ryom L, Monforte A, et al. Lack of association between use of efavirenz and death from suicide: evidence from the D:A:D study. *J Int AIDS Soc*. 2014;17(4 Suppl 3):19512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25394021>.
34. Strehlau R, Martens L, Coovadia A, et al. Absence seizures associated with efavirenz initiation. *Pediatr Infect Dis J*. 2011;30(11):1001-1003. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21633320>.
35. Teppler H, Brown DD, Leavitt RY, et al. Long-term safety from the raltegravir clinical development program. *Current HIV Res*. 2011;9(1):40-53. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21198432>.
36. Todd S, Rafferty P, Walker E, et al. Early clinical experience of dolutegravir in an HIV cohort in a larger teaching hospital. *Int J STD AIDS*. 2017;956462416688127.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28118801>.

37. van Dijk JH, Sutcliffe CG, Hamangaba F, Bositis C, Watson DC, Moss WJ. Effectiveness of efavirenz-based regimens in young HIV-infected children treated for tuberculosis: a treatment option for resource-limited settings. *PLoS One*. 2013;8(1):e55111. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23372824>.
38. Variava E, Sigauke FR, Norman J, et al. Brief report: late efavirenz-induced ataxia and encephalopathy: a case series. *J Acquir Immune Defic Syndr*. 2017;75(5):577-579. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28520619>.
39. Viswanathan P, Baro E, Soon G, Sherwat A, Struble KA. Neuropsychiatric adverse events associated with integrase strand transfer inhibitors. Presented at: Conference on Retroviruses and Opportunistic Infections. 2017. Seattle, Washington.
40. Walmsley S, Baumgarten A, Berenguer J, et al. Dolutegravir plus abacavir/lamivudine for the treatment of HIV-1 infection in antiretroviral therapy-naïve patients: week 96 and week 144 results from the SINGLE randomized clinical trial. *J Acquir Immune Defic Syndr*. 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26262777>.
41. Waters L, Fisher M, Winston A, et al. A Phase IV, double-blind, multicentre, randomized, placebo-controlled, pilot study to assess the feasibility of switching individuals receiving efavirenz with continuing central nervous system adverse events to etravirine. *AIDS*. 2011;25(1):65-71. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21099666>.
42. Arenas-Pinto A, Grund B, Sharma S, et al. Risk of suicidal behavior with use of efavirenz: results from the strategic timing of antiretroviral treatment trial. *Clin Infect Dis*. 2018;67(3):420-429. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29538636>.
43. Capetti AF, Di Giambenedetto S, Latini A, et al. Morning dosing for dolutegravir-related insomnia and sleep disorders. *HIV Med*. 2018;19(5):e62-e63. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28762661>.
44. Van de Wijer L, McHaile DN, de Mast Q, et al. Neuropsychiatric symptoms in Tanzanian HIV-infected children receiving long-term efavirenz treatment: a multicentre, cross-sectional, observational study. *Lancet HIV*. 2019;6(4):e250-e258. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30770324>.
45. Hammond CK, Eley B, Ing N, Wilmshurst JM. Neuropsychiatric and neurocognitive manifestations in HIV-infected children treated with efavirenz in South Africa—a retrospective case series. *Front Neurol*. 2019;10:742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31338063>.
46. Burman RJ, Wilmshurst JM, Gebauer S, Weise L, Walker KG, Donald KA. Seizures in children with HIV infection in South Africa: a retrospective case control study. *Seizure*. 2019;65:159-165. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30721872>.
47. Cuzin L, Pugliese P, Katlama C, et al. Integrase strand transfer inhibitors and neuropsychiatric adverse events in a large prospective cohort. *J Antimicrob Chemother*. 2019;74(3):754-760. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30534993>.
48. Hoffmann C, Llibre JM. Neuropsychiatric Adverse events with dolutegravir and other integrase strand transfer inhibitors. *AIDS Rev*. 2019;21(1):4-10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30899113>.